

Endocrine and Metabolic Manifestations of Invasive Fungal Infections and Systemic Antifungal Treatment

MICHAEL S. LIONAKIS, MD, ScD; GEORGE SAMONIS, MD; AND DIMITRIOS P. KONTOYIANNIS, MD, MS, ScD

Systemic fungal infections are increasingly reported in immunocompromised patients with hematological malignancies, recipients of bone marrow and solid organ allografts, and patients with AIDS. Mycoses may infiltrate endocrine organs and adversely affect their function or produce metabolic complications, such as hypopituitarism, hyperthyroidism or hypothyroidism, pancreatitis, hypoadrenalism, hypogonadism, hypernatremia or hyponatremia, and hypercalcemia. Antifungal agents used for prophylaxis and/or treatment of mycoses also have adverse endocrine and metabolic effects, including hypoadrenalism, hypogonadism, hypoglycemia, dyslipidemia, hypernatremia, hypocalcemia, hyperphosphatemia, hyperkalemia or hypokalemia, and hypomagnesemia. Herein, we review how mycoses and conventional systemic antifungal treatment can affect the endocrine system and cause metabolic abnormalities. If clinicians are equipped with better knowledge of the endocrine and metabolic complications of fungal infections and antifungal therapy, they can more readily recognize them and favorably affect outcome.

Mayo Clin Proc. 2008;83(9):1046-1060

ADH = antidiuretic hormone; AMB = amphotericin B; CYP = cytochrome P; DI = diabetes insipidus; LDL-C = low-density lipoprotein cholesterol; LH = luteinizing hormone; SIADH = syndrome of inappropriate ADH production

The incidence of systemic fungal infections is increasing in immunocompromised patients. Because these infections frequently disseminate, they may affect the function of the pituitary, thyroid, parathyroid, and adrenal glands; the pancreas; and the reproductive organs. Metabolic complications, such as electrolytic abnormalities, can also occur. Agents used for prophylaxis and/or treatment of mycoses also have adverse endocrine and metabolic effects (Table 1). We review how mycoses affect the endocrine system and cause metabolic derangements and discuss the endocrine and metabolic complications of antifungal agents. The focus is on only systemic antifungal agents for invasive mycoses, such as the polyenes (mainly amphotericin B [AMB] and its lipid formulations), the azoles (fluconazole, itraconazole, voriconazole, posaconazole,

and ketoconazole), and the echinocandins (caspofungin, micafungin, and anidulafungin). Agents that are used topically or have antifungal properties, such as pentamidine, trimethoprim-sulfamethoxazole, dapsone, and atovaquone, are not discussed.

We searched the MEDLINE and PubMed databases (from 1966 to February 2008) and abstracts from major infectious disease meetings held from 1992 to 2007 for published reports pertaining to the endocrine and metabolic manifestations of invasive mycoses and their antifungal treatment using terms such as *endocrine, metabolic, electrolytes, fungal, molds, yeast, aspergillosis, candida, pituitary, thyroid, pancreas, adrenal, ovaries, reproductive system, sodium, potassium, calcium, phosphorous, dyslipidemia, amphotericin B, azoles, and echinocandins*. Studies were selected on the basis of the importance of data (as determined by frequency of citations by subsequent publications in the specialty) and originality. Review articles by thought leaders in mycology and endocrinology were also included in our search.

PITUITARY INVOLVEMENT AND DYSFUNCTION

MYCOSES

In pituitary infections, which are rare, bacteria are the most common infecting microbes, whereas fungi are much less frequent.¹ Interestingly, the pituitary gland has a role in innate antifungal immunity; Breuel et al² showed that pituitary receptors sense circulating *Candida* glucans and respond by Toll-like receptor 4 (TLR4) and CD14 gene expression.

Pituitary involvement is also rare in patients with central nervous system fungal infections. Of 40 patients with intracranial fungal infections, only 6 (15%) had sellar distribution.³ Various mycoses can infiltrate the pituitary gland^{1,3-11} (Table 2). The pathogenesis of pituitary fungal infection involves (1) hematogenous spread, typically seen with disseminated opportunistic yeast or mold infections in immunosuppressed hosts; (2) extension from adjacent anatomical sites (ie, cryptococcal meningeal infection or aspergillosis of the sphenoid sinus, cavernous sinus, and skull base); or (3) iatrogenic inoculation (eg, with *Aspergillus* or *Candida* species) during transsphenoidal adenoma resection.^{1,3,4}

Pituitary fungal infections have no specific clinical features; like any pituitary process, they may cause headache, ophthalmoplegia due to optic chiasm compression, and/or

From the Department of Infectious Diseases, Infection Control and Employee Health, The University of Texas M. D. Anderson Cancer Center, Houston (M.S.L., D.P.K.); and University Hospital of Heraklion, University of Crete, Heraklion, Greece (G.S.)

Individual reprints of this article are not available. Address correspondence to Dimitrios P. Kontoyiannis, MD, MS, ScD, Department of Infectious Diseases, Infection Control and Employee Health, Unit 402, The University of Texas M. D. Anderson Cancer Center, 1515 Holcombe Blvd, Houston, TX 77030 (dkontoyi@mdanderson.org).

© 2008 Mayo Foundation for Medical Education and Research

TABLE 1. Adverse Endocrine and Metabolic Effects of Systemic Antifungal Agents

Antifungal agent	Endocrine effects	Metabolic effects
Polyenes		
Amphotericin B	Oligospermia in animals	Hypernatremia, hypokalemia, hyperkalemia, hypomagnesemia, hypocalcemia
Lipid formulations of amphotericin B	Pancreatitis	Hypernatremia, hypokalemia, hyperkalemia, hypomagnesemia, hypocalcemia, hyperphosphatemia, pseudohyperphosphatemia
Nystatin	Oligospermia in animals	Hypokalemia, hypomagnesemia, hypocalcemia
Azoles		
Ketoconazole	Hypothyroidism, Addison disease, decreased libido, impotence, gynecomastia, menstrual irregularities	Hypokalemia, hypocalcemia, hypoglycemia, dyslipidemia
Fluconazole	Pancreatitis, Addison disease, Conn-like syndrome, menstrual irregularities	Hypokalemia, hypoglycemia
Itraconazole	Pancreatitis, Addison disease, Conn-like syndrome, decreased libido, impotence, gynecomastia, menstrual irregularities	Hypokalemia, dyslipidemia
Voriconazole	Pancreatitis	Hypokalemia, hypomagnesemia, hypoglycemia, dyslipidemia
Posaconazole	Menstrual irregularities	None reported
Echinocandins		
Caspofungin, anidulafungin, micafungin	Pancreatitis	Hypokalemia, hypomagnesemia, hypercalcemia
Other systemic antimicrobial agents with antifungal properties		
Griseofulvin	Pancreatitis	None reported
Potassium iodide	Thyrototoxicosis (Jod-Basedow disease), hypothyroidism (Wolff-Chaikoff effect)	Hyperkalemia

endocrine dysfunction.^{1,3,4,11} Most such infections have no overt endocrine symptoms. However, hypopituitarism occurs rarely, manifesting most often as decreased libido and menstrual abnormalities because of luteinizing hormone (LH) and follicle-stimulating hormone insufficiency.^{11,12} Moreover, diabetes insipidus (DI) presenting as polydipsia and polyuria from impaired antidiuretic hormone (ADH) release is not uncommon.^{1,3,4,11,12} Indeed, DI occurs more frequently with pituitary infections than with adenomas, implying that posterior pituitary function is affected early during pituitary infections.^{1,12} Hyperprolactinemia and decreases in thyrotropin-releasing hormone and corticotropin also occur; conversely, endocrine function may be normal.^{1,3,4,11,12}

Fungal pituitary infections are radiographically indistinguishable from intrasellar bacterial infections and tumors. Consequently, they are often misdiagnosed as tumors, and diagnosis is made unexpectedly after surgery or at autopsy.^{1,4-11} Certain magnetic resonance imaging findings are thought to be specific for pituitary infection. Therefore, when peripheral enhancement, hypointensity, or calcifications are seen on T2-weighted images, infection is favored over a tumor.⁷ Investigators have used gallium-67 imaging for postoperative follow-up of patients with pituitary abscesses caused by *Aspergillus* spp, but more studies are needed to establish the role of imaging modalities in these

patients.¹³ The mortality rate with pituitary fungal infections exceeds 70% because of delayed recognition and the typically disseminated nature of these infections.^{1,10} Treatment comprises antifungal therapy and transsphenoidal resection, which is preferred over craniotomy to prevent intracranial dissemination.¹

Fungal infections can also influence sodium homeostasis by affecting pituitary ADH release. Hypernatremia secondary to DI or hyponatremia caused by the syndrome of inappropriate ADH production (SIADH) can develop in patients with mycoses. Diabetes insipidus occurs primarily with cryptococcal basilar meningitis¹⁴ but also with aspergillosis, candidiasis, zygomycosis, and blastomycosis.^{1,4,6,11,15,16} Fungal cryptococcal and coccidioidal meningitis rarely cause SIADH^{17,18}; the incidence of SIADH in patients with cryptococcal meningitis is approximately 8%.¹⁷ A well-recognized cause of SIADH is pneumonia.¹² Any fungal pneumonia can be complicated by SIADH. *Pneumocystis* pneumonia, caused by *Pneumocystis jiroveci* (formerly *Pneumocystis carinii*), is reportedly the most common cause of SIADH in patients with AIDS; antifungal treatment often corrects hyponatremia.¹⁹

ANTIFUNGAL AGENTS

Antidiuretic hormone acts by aquaporin 2–induced increase of the water permeability of renal collecting tubules.

TABLE 2. Mycoses Reported to Infiltrate Various Organs of the Endocrine System^a

Fungal infection	Pituitary gland	Thyroid gland	Pancreas	Adrenal gland	Male reproductive organs	Female reproductive organs
Aspergillosis	+	35.0%	4.7%	5.0%	+	+
Zygomycosis	—	3.0%	+	—	—	—
Fusariosis	—	—	+	—	+	—
Scedosporiosis	—	+	+	+	+	—
<i>Scopulariopsis</i> infection	+	+	—	—	—	—
Candidiasis	+	8.0%	2.9%	5.0%	—	+
Cryptococcosis	+	12.0%	3.8%	15.0%	+	+
Histoplasmosis	—	+	+	82.0%	+	+
Blastomycosis	—	+	+	10.0%	10.0%	+
Coccidioidomycosis	+	6.0%	+	36.0%	+	+
Paracoccidioidomycosis	+	4.0%	+	80.0%	+	+
Sporotrichosis	—	—	+	+	+	—
Trichosporonosis	—	+	+	—	—	+
Pneumocystosis	+	23.0% ^b	9.0%	15.0% ^b	—	—
Basidiobolomycosis	—	—	+	+	—	+
Chromoblastomycosis	—	—	+	—	—	—
<i>Saccharomyces</i> spp infection	—	—	+	—	—	—
<i>Malassezia</i> spp infection	—	—	+	+	—	—
<i>Trichophyton</i> spp infection	—	—	—	—	+	—

^a The percentages for postmortem infiltration of different endocrine organs correspond to the highest reported rate of infiltration in the literature. In cases in which infiltration is sporadic, percentages are not provided. + = sporadic infiltration; — = not reported in the literature.

^b Percentages refer to cases of extrapulmonary pneumocystosis.

Amphotericin B decreases aquaporin 2 expression in the kidney medulla, makes collecting tubules insensitive to ADH, and may rarely cause reversible nephrogenic DI.^{20,21} It also causes hypokalemia, a well-known etiology of nephrogenic DI; however, nephrogenic DI can occur without hypokalemia.^{20,21} Higher levels of prostaglandin E₂ and increased apoptosis seem to contribute to AMB-mediated DI.^{22,23} Amelioration of DI can be achieved with amiloride plus hydrochlorothiazide, indomethacin, and replacement of AMB-deoxycholate by lipid AMB formulations, which (for reasons that are not understood) cause DI less often.^{21,22,24}

THYROID INVOLVEMENT AND DYSFUNCTION

MYCOSES

The thyroid is resistant to microbial invasion because of its rich blood supply, iodine content, and capsule.²⁵ Although several fungi may infect the thyroid²⁵⁻⁴⁴ (Table 2), thyroid fungal infection occurs rarely and is clinically overt in a minority of patients.²⁵⁻²⁷ Of 415 previously reported cases of infectious thyroiditis (1900-1997), only 50 (12%) were fungal.²⁵⁻²⁷ The pathogenesis of fungal thyroiditis entails hematogenous or lymphatic spread.²⁵ Although most cases of thyroid fungal involvement occur during dissemination in immunosuppressed patients, isolated thyroid histoplasmosis, coccidioidomycosis, and pneumocystosis have been reported.^{38,40,44}

Aspergillus spp are the predominant causative fungus for thyroiditis and asymptomatic thyroid infiltration.²⁵⁻³¹

The literature contains 21 case reports of overt thyroiditis caused by *Aspergillus* spp, with thyroid infiltration found postmortem in 10% to 35% of aspergillosis cases.²⁵⁻³¹ In contrast, thyroid infiltration is less common in other mycoses^{9,25-32,36,45-49} (Table 2). Unsurprisingly, *P jiroveci* is the most common cause of fungal thyroiditis in patients with AIDS, reflecting the high incidence of pneumocystosis in these patients.^{12,44} Thyroiditis caused by *P jiroveci* is typically associated with aerosolized pentamidine prophylaxis, a known risk factor for extrapulmonary pneumocystosis.⁴⁴ In most cases, thyroid fungal involvement is an incidental finding at autopsy without antemortem evidence of thyroid dysfunction.^{12,25} Nevertheless, it can manifest as subacute thyroiditis with local and systemic symptoms.²⁵ Neck swelling and tenderness are common because of thyroid enlargement and dysphagia from esophageal compression; in severe cases, fatal respiratory failure from tracheal obstruction has occurred.^{25,28,50} Fungal thyroiditis typically begins with a brief hyperthyroid phase in which glandular destruction causes thyroid hormone release.^{12,25} Transient euthyroidism ensues, followed by hypothyroidism and, ultimately, recovery to euthyroidism.^{12,25} Hence, symptoms and laboratory evidence can range from those characteristic of hyperthyroidism (or even frank thyrotoxicosis) to those typical of hypothyroidism from fungal involvement, depending on the phase in which the condition is diagnosed.^{12,25} Hyperthyroidism is more common with aspergillosis and coccidioidomycosis, whereas hypothyroidism is more common with thyroiditis caused by *P jiroveci*.^{12,25-29,37,44} Whether this

pattern reflects the degree of glandular dysregulation by the aforementioned fungi or results from differences in the timing of diagnosis remains unknown.

Patients with acute severe fungal infections may develop euthyroid sick syndrome, characterized by decreased levels of triiodothyronine and increased levels of normal thyroxine and thyrotropin-releasing hormone, as described in cases of paracoccidioidomycosis.⁵¹ Because such laboratory findings signify diminished peripheral conversion of thyroxine to triiodothyronine, thyroid replacement therapy is unnecessary and detrimental.⁵² However, in one study, a decreased total triiodothyronine level was the best prognostic indicator of death from pneumonia caused by *P. jiroveci* in patients with AIDS.⁵³

Besides direct fungal invasion of the thyroid, mycotoxin secretion may influence thyroid function.^{54,55} Specifically, at doses relevant to human exposure levels, the *Aspergillus* toxin patulin decreased thyroxine levels in animals, and the *Fusarium* toxin deoxynivalenol increased them.^{54,55} Whether mycotoxin secretion during fungal infections affects thyroid function in humans remains unstudied.

ANTIFUNGAL AGENTS

In preclinical studies, imidazoles, such as ketoconazole, have had antithyroid effects because of interference with iodine and thyroid peroxidase.⁵⁶ Nevertheless, administration of 200 to 600 mg/d of ketoconazole for 1 month did not affect thyroid function in euthyroid, hyperthyroid, or hypothyroid patients.^{57,58} However, high-dose ketoconazole (1200 mg/d) may cause hypothyroidism.⁵⁹ Hypothyroidism also developed in 2 patients with chronic mucocutaneous candidiasis after prolonged (>3 months) low-dose (100-200 mg/d) treatment with ketoconazole.⁶⁰ Because patients with chronic mucocutaneous candidiasis frequently have hypothyroidism associated with adult polyglandular autoimmune syndrome type I, it remains unclear whether ketoconazole induced hypothyroidism in these patients.⁶⁰ No studies have implicated the triazoles in thyroid dysfunction.⁶¹

Use of potassium iodide is common in treating cutaneous sporotrichosis.⁶² Prolonged (>1 month) potassium iodide treatment is associated with reversible thyrotoxicosis in patients with coexistent "hot" thyroid nodules (Jod-Basedow disease) and with hypothyroidism in patients with excessive autoregulation (Wolff-Chaikoff effect).⁶²

CALCIUM AND PHOSPHORUS ABNORMALITIES AND PARATHYROID INVOLVEMENT

MYCOSES

Hypercalcemia is a well-known complication of many granulomatous disorders, including sarcoidosis, tuberculo-

sis, and lymphomas.⁶³ It is derived from extrarenal dysregulated production of 1,25-dihydroxyvitamin D due to interferon- γ -mediated expression of 1 α -hydroxylase by activated macrophages in granulomas.⁶⁴ Chronic fungal infections, including histoplasmosis, coccidioidomycosis, paracoccidioidomycosis, candidiasis, cryptococcosis, and pneumocystosis, can also cause granulomas and reversible hypercalcemia.⁶⁵⁻⁷⁰ Reports indicate that 3% of patients with various mycoses have hypercalcemia, whereas about 10% of patients with sarcoidosis do.⁶⁵⁻⁷⁰ Whether this reflects variation in granuloma formation or subtle differences in the mechanisms of hypercalcemia between these entities is unclear.

Studies have shown that the mechanisms of hypercalcemia differ with the various mycoses. For instance, coccidioidomycosis, the mycosis most commonly associated with hypercalcemia (incidence, 1.4%-25.0%), typically manifests late after initiation of antifungal treatment.^{67,68} Parathyroid hormone and parathyroid hormone-related peptide levels are normal. Hypercalcemia in coccidioidomycosis is not triggered by autonomous 1,25-dihydroxyvitamin D, prostaglandin, or osteoclast-activating factor production.^{67,68} Osseous coccidioidomycosis that causes bone resorption contributes in some cases, but hypercalcemia occurs even without bone involvement. Thus, the presence of an osteotropic substance resembling humoral hypercalcemia of malignancy has been postulated.^{67,68} Elevated levels of nephrogenous cyclic adenosine monophosphate, low levels of 1,25-dihydroxyvitamin D, and hypercalciuria with coccidioidomycosis have also been described in patients with hypercalcemia of malignancy.⁷¹

In contrast, studies have shown autonomous 1,25-dihydroxyvitamin D production with candidiasis, cryptococcosis, paracoccidioidomycosis, pneumocystosis, and histoplasmosis.^{65-67,69,70} Hypercalcemia has also been reported as part of the immune reconstitution syndrome in patients with AIDS and cryptococcosis after initiation of antiretroviral therapy.⁷² The underlying mechanism remains unclear, but it has been postulated that hypercalcemia is caused by restoration of the granulomatous host response, resulting in autonomous 1,25-dihydroxyvitamin D production after antiretroviral therapy.⁷³

Osteopontin, a glycoprotein ligand of the $\alpha_v\beta_3$ integrin that activates macrophages and osteoclasts,⁷⁴ is highly expressed by histiocytes in granulomas of diverse etiologies, including histoplasmosis.⁷⁵ Osteoclasts synthesize osteopontin and $\alpha_v\beta_3$ integrin, which both localize at bone resorption sites.⁷⁴ Hence, osteopontin may contribute to hypercalcemia via osteoclast activation and bone resorption; further studies are needed to elucidate its role in hypercalcemia associated with fungal infections. Another molecule implicated in osteoclast activation and bone resorption in

hypercalcemia of malignancy is the receptor activator of nuclear factor- κ B ligand; its role in hypercalcemia of mycoses also merits investigation.⁷⁶

Conversely, hypocalcemia and hyperphosphatemia caused by hypoparathyroidism have been described in a patient with pneumocystosis and parathyroid infiltration.⁷⁷ Supplementation with vitamin D and calcium was necessary to maintain eucalcemia.

ANTIFUNGAL AGENTS

Azoles inhibit ergosterol biosynthesis by blocking cytochrome P (CYP) 450-dependent 14 α -lanosterol demethylase.⁷⁸ The reported higher frequency of endocrine complications of ketoconazole than of other azoles derives from their different selectivity for fungal vs mammalian CYP450-dependent 14 α -lanosterol demethylase.⁷⁸

Ketoconazole inhibits *in vitro* conversion of 25-hydroxyvitamin D to 1,25-dihydroxyvitamin D by competitive inhibition of CYP450-dependent 1 α -hydroxylase.⁷⁹ Administration of 600 to 900 mg/d of ketoconazole has been shown to cause a rapid, dose-dependent decrease of 40% to 45% in 1,25-dihydroxyvitamin D levels in healthy participants and in patients with hyperparathyroidism, sarcoidosis, and tuberculosis, resulting in calcium-lowering in some, but not all, patients.⁸⁰⁻⁸² Conversely, the triazoles do not affect calcium homeostasis.

Polyene-induced hypomagnesemia resulting from iatrogenic Fanconi syndrome can cause hypocalcemia by adversely affecting parathyroid hormone secretion and action in the kidney.^{83,84} Cases of AMB- and nystatin-induced hypoparathyroidism and hypocalcemia have been reported.^{83,84} Conversely, caspofungin has been reported to induce hypercalcemia, but the mechanism by which it does so remains unknown.⁸⁵

Phosphate abnormalities are rarely associated with antifungal use. Specifically, because each 50-mg vial of liposomal AMB contains 37 mg of phosphorus, prolonged treatment with high-dose liposomal AMB can result in hyperphosphatemia, especially in patients with impaired renal function.⁸⁶ Nevertheless, because liposomal AMB can interfere with the Synchron LX-20 phosphorus assay (Beckman Coulter, Fullerton, CA), pseudohyperphosphatemia can occur, and caution is required before institution of phosphate-lowering therapy.⁸⁷

PANCREATIC INVOLVEMENT AND DYSFUNCTION

MYCOSES

Fungi may involve the pancreas in 2 patterns. The first is hematogenous involvement during dissemination^{9,34,45-49,88-98} (Table 2). Unless more than 80% of pancreatic tissue is affected, pancreatic insufficiency does not develop; thus,

involvement is asymptomatic and usually discovered postmortem. The second pattern is seeding by *Candida* spp of pancreatic tissue or pseudocysts after necrotizing pancreatitis.^{99,100} The incidence of such infection has increased considerably from about 7% of total postpancreatic infections in the 1980s to 40% in the 2000s after routine administration of prophylactic antibiotics in patients with necrotizing pancreatitis.^{99,100} Besides antibiotic prophylaxis that predisposes patients to fungal superinfection, the risk factors for fungal pancreatic infection are postpancreatitis abdominal surgical manipulation and the global immune dysregulation seen with necrotizing pancreatitis, as evidenced by decreases in CD4 lymphocytes and interleukin 2.⁹⁹⁻¹⁰¹

Most infections occur within 1 month after pancreatitis and derive from *Candida* spp translocated from the gut.^{99,100} Aspartic proteinase 6, secreted by *Candida* spp and regulated by the enhanced filamentous growth 1 (*EFG1*) transcription factor, has a role in enabling *Candida* spp to invade pancreatic tissue.¹⁰² In most studies, pancreatic candidal infection is associated with higher mortality rates than is bacterial pancreatic infection.⁹⁹ However, not all studies have shown such an association.⁹⁹ Interventions proposed to prevent pancreatic seeding by fungi in high-risk patients with necrotizing pancreatitis are selective bowel decontamination and fluconazole prophylaxis.^{99,100} The former approach is controversial; the latter decreases the incidence of such infections, but this reduction does not translate into survival benefits and carries the risk of selecting azole-resistant *Candida* spp.^{99,100}

ANTIFUNGAL AGENTS

Pancreatitis is rarely reported after treatment with griseofulvin, liposomal AMB (0.5%-1% of patients), micafungin, fluconazole, itraconazole, and voriconazole.¹⁰³⁻¹⁰⁷ Elevated levels of serum voriconazole can cause hypoglycemia, although the underlying mechanism is not understood and merits investigation.¹⁰⁷ Asian patients, who frequently have the CYP2C19 sequence variation that causes increased drug concentrations, may be especially predisposed to voriconazole-induced hypoglycemia.¹⁰⁸ Ketoconazole and fluconazole have also been associated with development of hypoglycemia.^{109,110} The mechanism for the former is unknown, whereas coadministration of fluconazole with oral hypoglycemic agents to patients with diabetes may inhibit the metabolism of the latter compounds and elevate their levels, thus predisposing patients to hypoglycemia.^{109,110} This interaction appears dose-dependent and agent-specific. Therefore, coadministration of 100 mg/d of fluconazole with tolbutamide or 200 mg/d of fluconazole with glimepiride or nateglinide increased the peak plasma concentrations and the area under the curve

of these compounds, hence heightening the risk of hypoglycemia.^{111,112} In contrast, coadministration of 50 mg/d of fluconazole with gliclazide or glibenclamide did not affect glycemic homeostasis.¹¹³ People with CYP2C9 sequence variations have a markedly diminished capacity to metabolize CYP2C9 substrates and might be particularly at risk for hypoglycemia in the setting of azole-hypoglycemic drug coadministration.¹¹⁴ However, more studies are needed to elucidate the frequency and relative risk of hypoglycemia as a result of different azole-hypoglycemic agent interactions.

ADRENAL INVOLVEMENT AND DYSFUNCTION

MYCOSES

The adrenal glands are the most common endocrine organs involved during infections, including mycoses. Frenkel¹¹⁵ suggested that excessive glucocorticoid levels in adrenal circulation (ie, 20-40 times higher than in the peripheral blood) generate local cell-mediated immunosuppression that predisposes patients to glandular invasion by microorganisms.

Isolated adrenal involvement has been reported in immunocompetent hosts with histoplasmosis, coccidioidomycosis, paracoccidioidomycosis, and pneumocystosis.¹¹⁶⁻¹¹⁹ Much more often, the adrenal glands are involved in widely disseminated mycoses; at autopsy, histoplasmosis and paracoccidioidomycosis are by far the most common fungal infections found to have infiltrated the adrenal glands^{9,34,45-49,92-95,98,120-123} (Table 2). Most such cases are asymptomatic, and antemortem diagnosis is rare. Adrenal function remains unaffected unless more than 90% of the adrenal cortex is obliterated, and so patients with disseminated mycoses may die of their infection before such destruction occurs.¹²⁰⁻¹²³ Also, manifestations of adrenal insufficiency are nonspecific and may be attributed erroneously to the mycosis.¹²⁰⁻¹²³

In addition to directly destroying adrenal tissue, host humoral factors generated during systemic infections, such as corticostatsins/defensins, may contribute to adrenal dysfunction. Specifically, these molecules inhibit adrenal steroidogenesis by interfering with corticotropin affinity at the receptor level; their role in affecting adrenal steroidogenesis in patients with systemic mycoses should be further explored.¹²⁴

Addison disease has been reported with histoplasmosis, coccidioidomycosis, paracoccidioidomycosis, cryptococcosis, blastomycosis, pneumocystosis, candidiasis, and infection with *Bipolaris* spp.^{118,120,122,123,125,126} Of these infections, histoplasmosis and paracoccidioidomycosis have the highest proclivity for the adrenal glands, and Addison disease occurs most frequently in patients with these infections (histoplasmosis, 7%-12%; paracoccidioidomycosis, 9%-14%).^{9,93,121} Prompt recognition and glucocorticoid

supplementation are fundamental for survival in such patients. In patients with blastomycosis and histoplasmosis, latency periods of several years between primary infection and adrenal insufficiency are not uncommon, signifying reactivation of quiescent adrenal foci.^{93,116,123}

Most adrenal insufficiency cases are irreversible and necessitate lifelong glucocorticoid supplementation. Nevertheless, recovery of adrenal function has been reported with antifungal treatment in histoplasmosis and paracoccidioidomycosis cases.^{127,128} Autopsy studies of paracoccidioidomycosis revealed fungal vasculitis in small arterioles, leading to tissue ischemia early during adrenal invasion.¹¹⁹ If left untreated, progression to glandular necrosis and irreversible endocrine impairment occur.¹¹⁹ Hence, prompt antifungal therapy may prevent irreversible pathological changes and salvage adrenal function.

Adrenal reserve can also be affected by systemic mycoses, such as histoplasmosis, cryptococcosis, coccidioidomycosis, paracoccidioidomycosis, pneumocystosis, and blastomycosis.^{93,116-118,120,123} In particular, 50% of patients with paracoccidioidomycosis have diminished adrenal reserve, as evidenced by attenuated cortisol increments after corticotropin administration.¹²⁹ Such patients are asymptomatic, but fatal adrenal failure can occur in conditions of physiological stress. Hence, when the condition of patients with mycoses unexpectedly deteriorates, the adrenal axis should be examined thoroughly.

Fungi might have differing affinities for infiltration of the different adrenal zones. For example, *Paracoccidioides* spp commonly infiltrate the zona reticularis, lowering dehydroepiandrosterone sulfate levels.^{129,130} Such a shift in adrenal steroidogenesis away from androgens and toward the essential glucocorticoid pathway is presumably an adaptive mechanism with infections.¹³⁰ Such patients may have impaired mineralocorticoid secretion with decreased aldosterone levels and elevated plasma renin activity because of zona glomerulosa infiltration. Mineralocorticoid deficiency can cause hyponatremia, hyperkalemia, and metabolic acidosis.¹²⁰⁻¹²³ In contrast, in patients with histoplasmosis, sparing of the zona glomerulosa and preservation of mineralocorticoid function is the rule.⁹³

Fungal adrenal involvement manifests radiographically as glandular enlargement with peripheral enhancement, central hypoattenuation, contour preservation, and calcifications during healing.¹³¹ Histoplasmosis causes bilateral symmetric adrenal enlargement, whereas paracoccidioidomycosis and blastomycosis cause asymmetric and occasionally unilateral involvement.¹³¹ These characteristics help differentiate fungal adrenal infections from metastatic disease, in which the adrenal contour is distorted, and from autoimmune adrenalitis, in which the glands are atrophic.¹³¹




FIGURE. The main pathways of steroidogenesis in the adrenal glands and inhibitory effects of ketoconazole. The principal enzymes inhibited by ketoconazole are shown in yellow, and the main steroid products whose synthesis is decreased by ketoconazole are shown in red. H = hydrogen; HSD = hydroxysteroid dehydrogenase; O = oxygen; OH = hydroxylase; OHD = hydroxy dehydrogenase; SCC = cholesterol side-chain cleavage.

Adapted from Hardman JG, Limbird LE, Gilman AG. *Goodman and Gilman's The Pharmacological Basis of Therapeutics*. 11th ed. New York, NY: McGraw-Hill, 2006, with permission of McGraw-Hill.

ANTIFUNGAL AGENTS

Azoles affect glucocorticoid and mineralocorticoid function by inhibiting CYP450-dependent enzymes involved in adrenal steroidogenesis. Specifically, ketoconazole is a dose-dependent reversible inhibitor of cholesterol desmolase, 17,20-lyase, 11β-hydroxylase, 17α-hydroxylase, and 18-hydroxylase (Figure).¹³²⁻¹³⁴ Critical factors that modulate ketoconazole's endocrine effects are dosage and timing of administration because the kinetics of adrenal inhibition reflect serum ketoconazole concentrations.¹³²⁻¹³⁵ Thus, no effect on glucocorticoid synthesis is seen after administration of 200 mg of ketoconazole.^{135,136} However, use of

400 or 600 mg of ketoconazole transiently blunts plasma cortisol response to corticotropin without affecting basal cortisol or pituitary corticotropin secretion.¹³⁴⁻¹³⁷ This effect lasts 8 hours but normalizes by 16 hours after ketoconazole administration.¹³⁴⁻¹³⁷ Higher doses produce more pronounced adrenal dysfunction. Nevertheless, overt adrenal insufficiency is relatively infrequent with ketoconazole, presumably because of a compensatory increase in corticotropin. However, overt adrenal insufficiency can occur with administration of 600 to 1200 mg/d of ketoconazole in divided doses; as opposed to once-daily ketoconazole use, such regimens produce sustained adrenal inhibition with-

out allowing for compensatory normalization of adrenal steroidogenesis for a substantial part of the day.¹³⁴⁻¹³⁸ In contrast, adrenal crisis has occurred with ketoconazole dosages as low as 400 mg/d, suggesting host-specific genetic susceptibility to adrenal dysfunction.¹³⁹

In addition to inhibiting glucocorticoid synthesis, ketoconazole is a dose-dependent, reversible, competitive antagonist at the glucocorticoid receptor level.¹⁴⁰ In fact, because of its antiglucocorticoid properties and rapid action onset, physicians have used ketoconazole (800-1200 mg/d) in palliative treatment of Cushing disease, adrenal tumors, and ectopic corticotropin production by small-cell lung carcinoma or carcinoid tumors.¹⁴¹⁻¹⁴³ The aforementioned interaction of ketoconazole with the glucocorticoid receptor is restricted to imidazoles and does not occur with triazoles.¹⁴⁴

Clinicians have used ketoconazole's ability to inhibit mineralocorticoid synthesis for palliative treatment of primary hyperaldosteronism due to adrenal adenomas or hyperplasia.^{135,145} However, the accumulation of 11-deoxycortisol, corticosterone, and 11-deoxycorticosterone induced by inhibition of cortisol and aldosterone synthesis (Figure) can cause hypersensitivity to the mineralocorticoid properties of these compounds; indeed, hypertension develops in 14% to 27% of patients after long-term high-dose ketoconazole use; hypokalemia and edema may also develop, suggesting ketoconazole-induced Conn syndrome.¹⁴⁶

Fluconazole inhibits CYP450-dependent enzymes in cell cultures and animals, but concentrations required for these effects far exceed its therapeutic range. Consequently, adrenal function is unaffected in patients receiving 400 mg/d of fluconazole.^{147,148} However, the literature contains several reports of fluconazole-induced reversible Addison disease occurring within 24 to 48 hours of treatment.^{147,149} Critically ill patients already at increased risk of adrenal insufficiency who receive high-dose fluconazole (800 mg/d) are most predisposed to developing adrenal insufficiency; however, it has been reported even with dosing of 200 mg/d.¹⁴⁹ Fluconazole has reversed hypercortisolism in adrenal carcinoma-induced Cushing syndrome cases.¹⁵⁰

Although itraconazole does not inhibit adrenal steroidogenesis at 100 to 400 mg/d,¹⁵¹ reversible adrenal insufficiency has been reported at 600 mg/d.¹⁵² Also, combination of itraconazole with high-potency inhaled glucocorticoids can suppress the hypothalamic-pituitary-adrenal axis.¹⁵³ Such suppression occurs because itraconazole, an inhibitor of CYP3A4, decreases glucocorticoid clearance and increases serum glucocorticoid levels.¹⁵⁴ In a study of 25 patients receiving itraconazole and inhaled budesonide, adrenal insufficiency developed in 11 patients and Cushing syndrome in 1 patient.¹⁵³ Adrenal function recovered in only 1 patient, whereas the remaining 10 had persistent adrenal insufficiency 2 to 10 months after treat-

ment discontinuation.¹⁵³ Variation in CYP3A4 activity or glucocorticoid receptor sequence variations may explain such host-specific phenotypic diversity in adrenal function manifestations.¹⁵⁵ In addition, mineralocorticoid-induced edema, hypokalemia, and hypertension have been reported in 63% of patients receiving high-dose itraconazole (600 mg/d).¹⁵² No adrenal effects have been reported with voriconazole or posaconazole.

POTASSIUM AND MAGNESIUM ABNORMALITIES INDUCED BY ANTIFUNGAL AGENTS

Hypokalemia develops in up to 90% of AMB-deoxycholate-treated patients and may occur and persist even weeks after treatment discontinuation.¹⁵⁶ Amphotericin B forms intramembranous pores in distal convoluted renal tubules, increasing their permeability and potassium urinary wasting.¹⁵⁶ Apoptosis is an important modulator of this process.²³ Additionally, AMB causes renal tubular acidosis type I because of a defect in distal tubule hydrogen potassium adenosine triphosphatase and promotes potassium urinary loss.¹⁵⁶ Moreover, AMB enhances sodium reabsorption in the distal colon, causing fecal potassium loss.¹⁵⁶ A recent study suggested that preexisting proteinuria may protect from AMB-deoxycholate-induced hypokalemia because binding of AMB to albumin in the tubular lumen may decrease the free drug fraction available to bind to the luminal epithelial surface.¹⁵⁷

Reversible, dose-dependent hypokalemia is more often caused by AMB-deoxycholate than by lipid AMB formulations¹⁵⁶; the latter have less affinity for mammalian cell membranes, thus reducing the incidence of nephrotoxicity. However, dose-dependent liposomal AMB-induced hypokalemia does occur; a recent study showed a 30% and 16% incidence of hypokalemia (potassium level <3 mEq/L [to convert to mmol/L, multiply by 1.0]) in patients treated with 10 vs 3 mg/kg of liposomal AMB daily, respectively.¹⁵⁸ Hypokalemia can be ameliorated by administration of amiloride or spironolactone or by dilution of AMB in fat emulsion instead of dextrose.¹⁵⁹⁻¹⁶¹ Nystatin causes hypokalemia by a similar mechanism.⁸⁴ Fluconazole, itraconazole, and voriconazole also cause hypokalemia in up to 9%, 12%, and 16% of treated patients, respectively.^{162,163} Ventricular fibrillation and rhabdomyolysis have been reported with severe itraconazole-mediated hypokalemia.¹⁶⁴ Echinocandins cause hypokalemia in 2% to 11% of patients, and studies have suggested a dose-dependent hypokalemic effect for caspofungin.^{85,165} The precise mechanisms by which azoles and echinocandins cause hypokalemia have not been elucidated.

The AMB lipid complex and AMB-deoxycholate are also associated with hyperkalemia via intracellular potassium

shift.¹⁶⁶ Renal insufficiency and rapid infusion (<1 hour) increase AMB serum levels and the risk of hyperkalemia and ventricular fibrillation¹⁶⁶; this increased risk can be avoided by slower infusion (4–6 hours) and AMB administration during hemodialysis.¹⁶⁶ Renal failure also predisposes potassium iodide-treated patients to hyperkalemia.⁶²

Hypomagnesemia occurs in 30% to 75% of AMB-treated patients because of increased urinary magnesium secretion from AMB-induced tubular defects in distal convoluted tubules.^{83,167} The incidence of hypomagnesemia increases during the second week of AMB administration, peaks by the fourth week, and then plateaus.¹⁶⁷ Hypomagnesemia, which is dose-dependent and reversible, is more common with AMB-deoxycholate than with lipid AMB formulations; it may persist for weeks after AMB discontinuation.¹⁶⁷ Amiloride ameliorated AMB-mediated hypomagnesemia in some studies.¹⁵⁹ Up to 63% of patients treated with voriconazole and up to 3% of those treated with caspofungin have been shown to develop hypomagnesemia.^{85,163,165}

EFFECTS OF ANTIFUNGAL AGENTS ON LIPID METABOLISM

Ketoconazole inhibits cholesterol synthesis in a dose-dependent fashion by blocking conversion of lanosterol to cholesterol.^{78,168} Other lipid-modifying properties of ketoconazole include inhibition of macrophage-induced low-density lipoprotein cholesterol (LDL-C) oxidation, decreased lipoprotein lipase and 3-hydroxy-3-methylglutaryl coenzyme A reductase activities, inhibition of intestinal cholesterol absorption and bile acid synthesis, and upregulation of LDL-C receptor activity.^{169–171}

At dosages of 200, 400, 800, and 1200 mg/d, ketoconazole reduced LDL-C levels by 10%, 13%, 33%, and 41% and reduced total cholesterol levels by 7%, 10%, 19%, and 27%, respectively, without affecting high-density lipoprotein or very-low-density-lipoprotein levels but with concomitant hypertriglyceridemia.^{78,168,172} Cholesterol reduction occurs rapidly (within 1 month) but is transient because cholesterol levels increase toward pretreatment levels despite ketoconazole continuation.¹⁶⁸ Conversely, hypercholesterolemia and hypertriglyceridemia have been reported after treatment with itraconazole and voriconazole, but the underlying mechanism remains unclear.^{163,173}

INVOLVEMENT AND DYSFUNCTION OF THE MALE REPRODUCTIVE SYSTEM

Mycoses

Fungal epididymo-orchitis can occur as an isolated entity or, more often, during disseminated infection.¹⁷⁴ Blastomycosis is the predominant mycosis involving the male repro-

ductive organs in up to 10% of cases, although other fungal infections have been reported^{45–49,91,95,174–178} (Table 2).

Fungal epididymo-orchitis may be asymptomatic and discovered postmortem or can cause scrotal swelling and pain, requiring antifungal therapy and surgical resection.¹⁷⁴ As with any gonadal infection, fungal epididymo-orchitis can cause infertility because of gonadal destruction and resultant azoospermia.¹⁷⁹ In addition to invading tissue, fungi can contribute to infertility by inducing antisperm effects and secreting mycotoxin. *Candida guilliermondii* and *Candida albicans* are able to inhibit sperm viability and motility in vitro.^{180–182} In fact, a proportion of infertile men and women have antibodies positive for *C. guilliermondii*.^{180,181} The meaning of this finding is unknown, but restoration of fertility was achieved in some patients after eradication of *C. guilliermondii* by ketoconazole.^{180,181} *Cryptococcus neoformans* can also cause hypospermia and teratospermia.¹⁸³ The *Fusarium* toxin zearalenone and its metabolite zearalenol bind as agonists to estrogen receptor- α and - β , causing hyperestrogenism-mediated decreases in testosterone and libido, azoospermia, and feminization in mammals.¹⁸⁴ Whether such hyperestrogenic effects occur in patients with fusariosis is unclear.

ANTIFUNGAL AGENTS

Ketoconazole inhibits testosterone production by Leydig cells in vitro and impairs fertility in animals.¹⁸⁵ The most sensitive site of ketoconazole action in male gonads is 17,20-lyase, which is involved in sex-hormone production (Figure)^{134,136,186}; a single 200-mg dose has been reported to selectively inhibit testosterone without altering cortisol levels.^{134,136,186} Testosterone inhibition is reversible and dose-dependent, and an inverse relationship exists between serum ketoconazole and testosterone concentrations.^{134,136,137,186} Administration of 200 to 600 mg of ketoconazole lowers testosterone levels within 2 to 4 hours, but they recover within 24 hours.^{137,186} A lower nadir and longer duration of suppressed testosterone levels occur with increasing ketoconazole doses; in a substantial portion of patients, testosterone suppression was sustained throughout the day with 800 mg of ketoconazole.^{137,186}

These antiandrogenic properties account for hypogonadism during high-dose ketoconazole treatment.¹⁸⁶ In patients receiving more than 800 mg/d, oligospermia and azoospermia have been reported in 66% and 22% of patients, respectively.^{12,134,172} Decreased libido and impotence developed in 2% to 4% of patients at 400 mg/d and in 12% to 14% at 800 to 1200 mg/d.^{12,134,172} Finally, reversible gynecomastia appeared after as little as 1 month or after as many as 32 months of ketoconazole treatment in 2% to 7%, 14%, 23%, and 33% of patients treated with 400, 800, 1200, and 2000 mg/d, respectively.^{12,134,172} Ketoconazole-mediated in-

creases in the estradiol:testosterone ratio appear to be fundamental to the pathogenesis of gynecomastia.¹⁸⁷

Ketoconazole's antiandrogenic properties have clinical applications. For instance, Glass¹⁸⁸ advocated ketoconazole for testing gonadotropin reserve in infertile men. Because of its rapid onset of action and ability to block both adrenal and gonadal androgen synthesis, ketoconazole is used for advanced prostate cancer.¹⁸⁹ Administration of 400 mg every 8 hours causes initial testosterone lowering to castration levels.^{134,189} Despite subsequent testosterone-level increases driven by compensatory increases in LH levels, many patients with prostate cancer have striking biochemical and clinical responses, leading to tumor regression and symptom improvement. To prevent compensatory increases in LH levels and diurnal testosterone fluctuations observed with ketoconazole monotherapy, a gonadotropin-releasing hormone analogue such as leuprolide is often coadministered.^{134,189}

Ketoconazole has also been used for treatment of autonomous Leydig cell hyperactivity in children with precocious puberty, resulting in remarkable behavioral and clinical improvement with decreased growth and skeletal maturation rates.¹⁹⁰ Topical ketoconazole has been suggested as an adjunct to finasteride for androgenetic alopecia.¹⁹¹ By inhibiting 5 α -reductase and competitive binding to androgen receptors,^{192,193} ketoconazole may prevent dihydrotestosterone attachment to androgen receptors in hair follicles, a key step in alopecia development.

At extremely high doses, fluconazole has caused mild testosterone inhibition in vitro, exhibiting less than 1% of ketoconazole's potency and causing reversible oligospermia and sperm dysmotility in rabbits.¹⁹⁴ However, administration at 50 mg/d does not lower basal or gonadotropin-stimulated testosterone levels, and no antiandrogenic complications have been observed in patients treated at even higher doses.¹⁹⁵ Similarly, at 200 to 400 mg/d, itraconazole does not affect testosterone production, and decreased libido, impotence, and gynecomastia are sporadically reported in less than 1% of treated patients.^{151,173} In animals, AMB and nystatin cause oligospermia and sperm dysmotility, but their effects on human fertility remain unknown.¹⁹⁶

INVOLVEMENT AND DYSFUNCTION OF THE FEMALE REPRODUCTIVE SYSTEM

MYCOSES

Several mycoses can infiltrate the uterus, salpinx, and ovaries during dissemination^{45,93,95,174,197-203} (Table 2). Conversely, isolated genital blastomycosis and histoplasmosis can be acquired by sexual transmission.^{199,202} Fungal infections of female reproductive organs can be asymptomatic or present as endometritis and/or tubo-ovarian abscess,

causing menstrual irregularities, menorrhagia, dysmenorrhea, anovulation, and/or infertility.¹⁹⁷⁻²⁰³

Mycoses can cause dysregulation of the reproductive endocrine axis via mechanisms other than tissue invasion. For example, Arora et al²⁰⁴ reported hyperprolactinemia-induced amenorrhea and galactorrhea in a woman with blastomycosis involving the pleural cavity. This was attributed to prolactin production by the chest-wall inflammatory process, a well-described condition that causes prolactin release. Furthermore, the *Fusarium* toxin zearalenone can induce hyperestrogenism in female mammals but rarely in humans, leading to decreased LH and progesterone levels, infertility, vulvovaginitis, and decreased milk production.¹⁸⁴

ANTIFUNGAL AGENTS

Ketoconazole inhibits ovarian aromatase in vitro and in animals, but aromatase blockade in human ovaries is less pronounced than androgen blockade by 17,20-lyase.²⁰⁵ Consequently, ketoconazole does not cause sustained or predictable estrogen-level lowering in women; a single 200-mg dose caused a 16% decrease in estradiol levels at the luteal phase.^{134,205,206} However, such modest decreases are transient, and increases to pretreatment levels occur with continuous treatment. In contrast, women with adrenal tumor-induced estrogen oversecretion have experienced marked, sustained estradiol reductions with high-dose ketoconazole.¹⁴² However, it is unclear whether estrogen reduction resulted from aromatase inhibition or androgen synthesis blockade and therefore estrogen precursor deprivation. Ketoconazole also decreases ovarian progesterone production, impairs early uterine implantation, and decreases decidual response after implantation, suggesting interference with early pregnancy.²⁰⁷ Ketoconazole-mediated sex-hormone dysregulation has been reported to result in dose-dependent menstrual irregularities in 6% and 19% of women given 400 and 800 mg/d, respectively.^{172,208}

Because of its potential to inhibit ovarian aromatase, high-dose ketoconazole has been studied in women with advanced breast cancer resistant to conventional chemotherapy.^{59,206} Ketoconazole did not reliably reduce estrogen levels, and only 1 of 14 patients had a partial response after adding ketoconazole to aminoglutethimide.²⁰⁶ Conversely, because of its ability to substantially block ovarian androgen synthesis, physicians have used ketoconazole for ovarian hyperandrogenism syndromes, including polycystic ovarian syndrome and hyperthecosis, with considerable improvement in acne, hirsutism, and amenorrhea.²⁰⁹

Triazoles do not appreciably affect production of sex hormones in women, and menstrual irregularities are rarely reported with fluconazole and itraconazole; however, 6% of patients treated with posaconazole for more than 6 months developed such symptoms.^{173,210} Because flucon-

azole inhibits estrogen metabolism and increases ethinyl estradiol levels in women taking oral contraceptives,²¹¹ long-term fluconazole therapy in these women may induce hyperestrogenic effects. Such interaction with oral contraceptive treatment may also occur with voriconazole.²¹² Conversely, inadvertent pregnancies have been reported in patients taking oral contraceptives and itraconazole because estrogen metabolism was enhanced by the latter.²¹³

CONCLUSION

Fungal infections may have pleiotropic effects on the endocrine system associated with pituitary, thyroid, parathyroid, pancreatic, adrenal, and reproductive organ infiltration and may lead to metabolic and electrolytic disturbances. Similar complications may arise during treatment of such infections with systemic antifungal agents. A better understanding and high index of suspicion for these infections might result in earlier recognition and in amelioration of the morbidity and mortality associated with the complex interaction between fungi and/or their treatment and the endocrine/metabolic system.

REFERENCES

- Mielke B, Weir B, Oldring D, von Westarp C. Fungal aneurysm: case report and review of the literature. *Neurosurgery*. 1981;9(5):578-582.
- Breuel KF, Kougias P, Rice PJ, et al. Anterior pituitary cells express pattern recognition receptors for fungal glucans: implications for neuroendocrine immune involvement in response to fungal infections. *Neuroimmunomodulation*. 2004;11(1):1-9.
- Dubey A, Patwardhan RV, Sampth S, Santosh V, Kolluri S, Nanda A. Intracranial fungal granuloma: analysis of 40 patients and review of the literature. *Surg Neurol*. 2005;63(3):254-260.
- Pinzer T, Reiss M, Bourquain H, Krishnan KG, Schackert G. Primary aspergillosis of the sphenoid sinus with pituitary invasion—a rare differential diagnosis of sellar lesions. *Acta Neurochir (Wien)*. 2006;148(10):1085-1090.
- Yeung IY, Whitelaw BC, Hortobágyi T, et al. Phaeohyphomycosis: an unusual pituitary mass. *Br J Neurosurg*. 2005;19(5):442-445.
- Heary RF, Maniker AH, Wolansky LJ. Caudal pituitary abscess: case report. *Neurosurgery*. 1995;36(5):1009-1012.
- Yu YQ, Jiang XX, Gao YJ. MRI of a pituitary cryptococcoma simulating an adenoma. *Neuroradiology*. 1995;37(6):449-450.
- Scanarini M, Rotilio A, Rigobello L, Pomes A, Parenti A, Alessio L. Primary intrasellar coccidioidomycosis simulating a pituitary adenoma. *Neurosurgery*. 1991;28(5):748-751.
- Murray HW, Littman ML, Roberts RB. Disseminated paracoccidioidomycosis (South American blastomycosis) in the United States. *Am J Med*. 1974;56(2):209-220.
- Bartlett JA, Hulette C. Central nervous system pneumocystosis in a patient with AIDS. *Clin Infect Dis*. 1997;25(1):82-85.
- Stalldecker G, Molina HA, Antelo N, Arakaki T, Sica RE, Basso A. Hypopituitarism caused by colonic carcinoma metastasis associated with hypophysial aspergillosis [in Spanish]. *Medicina (B Aires)*. 1994;54(3):248-252.
- Aron DC. Endocrine complications of the acquired immunodeficiency syndrome. *Arch Intern Med*. 1989;149(2):330-333.
- Parker KM, Nicholson JK, Cezayirli RC, Biggs PJ. Aspergillosis of the sphenoid sinus: presentation as a pituitary mass and postoperative gallium-67 imaging. *Surg Neurol*. 1996;45(4):354-358.
- Juffermans NP, Verbon A, Van der Poll T. Diabetes insipidus as a complication of cryptococcal meningitis in an HIV-infected patient. *Scand J Infect Dis*. 2002;34(5):397-398.
- Ryan M, Yeo S, Maguire A, et al. Rhinocerebral zygomycosis in childhood acute lymphoblastic leukaemia. *Eur J Pediatr*. 2001;160(4):235-238.
- Kelly PM. Systemic blastomycosis with associated diabetes insipidus. *Ann Intern Med*. 1982;96(1):66-67.
- Stockstill MT, Kauffman CA. Comparison of cryptococcal and tuberculous meningitis. *Arch Neurol*. 1983;40(2):81-85.
- Webb M, Ziauddin A, Okusa MD. Coccidioidomycosis meningitis and syndrome of inappropriate antidiuretic hormone. *Am J Med Sci*. 2002;324(3):155-157.
- Tang WW, Kaptein EM, Feinstein EI, Massry SG. Hyponatremia in hospitalized patients with the acquired immunodeficiency syndrome (AIDS) and the AIDS-related complex. *Am J Med*. 1993;94(2):169-174.
- Kim SW, Yeum CH, Kim S, Oh Y, Choi KC, Lee J. Amphotericin B decreases adenylyl cyclase activity and aquaporin-2 expression in rat kidney. *J Lab Clin Med*. 2001;138(4):243-249.
- Späth-Schwalbe E, Koschuth A, Dietzmann A, Schanz J, Possinger K. Successful use of liposomal amphotericin B in a case of amphotericin B-induced nephrogenic diabetes insipidus. *Clin Infect Dis*. 1999;28(3):680-681.
- Höhler T, Teuber G, Wanitschke R, Meyer zum Büschenfeld KH. Indomethacin treatment in amphotericin B induced nephrogenic diabetes insipidus. *Clin Invest*. 1994;72(10):769-771.
- Varlam DE, Siddiq MM, Parton LA, Rüßmann H. Apoptosis contributes to amphotericin B-induced nephrotoxicity. *Antimicrob Agents Chemother*. 2001;45(3):679-685.
- Kirchlechner V, Koller DY, Seidl R, Waldhauser F. Treatment of nephrogenic diabetes insipidus with hydrochlorothiazide and amiloride. *Arch Dis Child*. 1999;80(6):548-552.
- Berger SA, Zonszein J, Villamena P, Mittman N. Infectious diseases of the thyroid gland. *Rev Infect Dis*. 1983;5(1):108-122.
- Goldani LZ, Zavascki AP, Maia AL. Fungal thyroiditis: an overview. *Mycopathologia*. 2006;161(3):129-139.
- Yu EH, Ko WC, Chuang YC, Wu TJ. Suppurative *Acinetobacter baumannii* thyroiditis with bacteremic pneumonia: case report and review. *Clin Infect Dis*. 1998;27(5):1286-1290.
- Sion ML, Armenaka MC, Georgiadis I, Paraskevopoulos G, Nikolaidis I. *Aspergillus fumigatus* abscesses of the thyroid with obstruction of the esophagus. *Thyroid*. 2004;14(9):786-788.
- Winzelberg GG, Gore J, Yu D, Vagenakis AG, Braverman LE. *Aspergillus flavus* as a cause of thyroiditis in an immunosuppressed host. *Johns Hopkins Med J*. 1979;144(3):90-93.
- Kalina PH, Campbell RJ. *Aspergillus terreus* endophthalmitis in a patient with chronic lymphocytic leukemia. *Arch Ophthalmol*. 1991;109(1):102-103.
- Iwen PC, Rupp ME, Bishop MR, et al. Disseminated aspergillosis caused by *Aspergillus ustus* in a patient following allogeneic peripheral stem cell transplantation. *J Clin Microbiol*. 1998;36(12):3713-3717.
- Mori T, Egashira M, Kawamata N, et al. Zygomycosis: two case reports and review of reported cases in the literature in Japan [in Japanese]. *Nippon Ishinkin Gakkai Zasshi*. 2003;44(3):163-179.
- Riddell J IV, Chenoweth CE, Kauffman CA. Disseminated *Scedosporium apiospermum* infection in a previously healthy woman with HELLP syndrome. *Mycoses*. 2004;47(9-10):442-446.
- Berenguer J, Rodríguez-Tudela JL, Richard C, et al. Deep infections caused by *Scedosporium prolificans*: a report on 16 cases in Spain and a review of the literature. *Medicine (Baltimore)*. 1997;76(4):256-265.
- Wuyts WA, Molzahn H, Maertens J, et al. Fatal *Scopulariopsis* infection in a lung transplant recipient: a case report. *J Heart Lung Transplant*. 2005 Dec;24(12):2301-2304. Epub 2005 Sep 28.
- Gandhi RT, Tollin SR, Seely EW. Diagnosis of *Candida* thyroiditis by fine needle aspiration. *J Infect*. 1994;28(1):77-81.
- Avram AM, Sturm CA, Michael CW, Sisson JC, Jaffe CA. Cryptococcal thyroiditis and hyperthyroidism. *Thyroid*. 2004;14(6):471-474.
- Goldani LZ, Klock C, Diehl A, Monteiro AC, Maia AL. Histoplasmosis of the thyroid. *J Clin Microbiol*. 2000;38(10):3890-3891.
- Babu A, Lacuesta E, Patel S. Cervical blastomycosis masquerading as a thyroid mass. *Endocr Pract*. 2005;11(1):65-68.
- Smilack JD, Argueta R. Coccidioidal infection of the thyroid. *Arch Intern Med*. 1998;158(1):89-92.
- Ito T, Ishikawa Y, Fujii R, et al. Disseminated *Trichosporon capitatum* infection in a patient with acute leukemia. *Cancer*. 1988;61(3):585-588.
- Gold JW, Poston W, Mertelsmann R, et al. Systemic infection with *Trichosporon cutaneum* in a patient with acute leukemia: report of a case. *Cancer*. 1981;48(10):2163-2167.

43. Koyanagi T, Nishida N, Osabe S et al. Autopsy case of disseminated *Trichosporon inkin* infection identified with molecular biological and biochemical methods. *Pathol Int*. 2006;56(12):738-743.
44. Guttler R, Singer PA, Axline SG, Greaves TS, McGill JJ. *Pneumocystis carinii* thyroiditis: report of three cases and review of the literature. *Arch Intern Med*. 1993;153(3):393-396.
45. Bodey G, Bueltmann B, Duguid W, et al. Fungal infections in cancer patients: an international autopsy survey. *Eur J Clin Microbiol Infect Dis*. 1992;11(2):99-109.
46. Groll AH, Shah PM, Mentzel C, Schneider M, Just-Nuebling G, Huebner K. Trends in the postmortem epidemiology of invasive fungal infections at a university hospital. *J Infect*. 1996;33(1):23-32.
47. Antinori S, Galimberti L, Magni C, et al. *Cryptococcus neoformans* infection in a cohort of Italian AIDS patients: natural history, early prognostic parameters, and autopsy findings. *Eur J Clin Microbiol Infect Dis*. 2001;20(10):711-717.
48. Ng VL, Yajko DM, Hadley WK. Extrapulmonary pneumocystosis. *Clin Microbiol Rev*. 1997;10(3):401-418.
49. Huntington RW, Waldmann WJ, Sargent JA, O'Connell H, Wybel R, Croll D. Pathological and clinical observations on 142 cases of fatal coccidioidomycosis with necropsy. In: Ajella L, ed. *Coccidioidomycosis*. Tucson, AZ: University of Arizona Press;1967:143-167.
50. Kishi Y, Negishi M, Kami M, et al. Fatal airway obstruction caused by invasive aspergillosis of the thyroid gland. *Leuk Lymphoma*. 2002;43(3):669-671.
51. Kiy Y, Machado JM, Mendes RP, Barraviera B, Pereira PC, Cury PR. Paracoccidioidomycosis in the region of Botucatu (state of São Paulo, Brazil): evaluation of serum thyroxine (T4) and triiodothyronine (T3) levels and of the response to thyrotropin releasing hormone (TRH). *Mycopathologia*. 1988;103(1):3-9.
52. Brent GA, Hershman JM. Thyroxine therapy in patients with severe nonthyroidal illnesses and low serum thyroxine concentration. *J Clin Endocrinol Metab*. 1986;63(1):1-8.
53. Fried JC, LoPresti JS, Micon M, Bauer M, Tuchschildt JA, Nicoloff JT. Serum triiodothyronine values: prognostic indicators of acute mortality due to *Pneumocystis carinii* pneumonia associated with the acquired immunodeficiency syndrome. *Arch Intern Med*. 1990;150(2):406-409.
54. Selmanoglu G, Kockaya EA. Investigation of the effects of patulin on thyroid and testis, and hormone levels in growing male rats. *Food Chem Toxicol*. 2004;42(5):721-727.
55. Rotter BA, Thompson BK, Lessard M, Trenholm HL, Tryphonas H. Influence of low-level exposure to Fusarium mycotoxins on selected immunological and hematological parameters in young swine. *Fundam Appl Toxicol*. 1994;23(1):117-124.
56. Comby F, Lagorce JF, Buxeraud J, Raby C. Antithyroid action of ketoconazole: in-vitro studies and rat in-vivo studies. *J Pharm Pharmacol*. 1994;46(1):50-53.
57. De Pedrini P, Montemurro G, Tommaselli A, Costa C. No effect of ketoconazole on thyroid function of normals and thyrotoxic patients. *Int J Tissue React*. 1988;10(3):189-192.
58. De Pedrini P, Tommaselli A, Montemurro G. No effect of ketoconazole on thyroid function of normal subjects and hypothyroid patients. *Int J Clin Pharmacol Res*. 1988;8(6):485-488.
59. Namer M, Khater R, Frenay M, Boulblil JL. High-dose ketoconazole in the treatment of advanced cancer of the breast [in French] [letter]. *Bull Cancer*. 1986;73(1):89.
60. Kitching NH. Hypothyroidism after treatment with ketoconazole [letter]. *Br Med J*. 1986;293(6553):993-994.
61. Devenport MH, Crook D, Wynn V, Lees LJ. Metabolic effects of low-dose fluconazole in healthy female users and non-users of oral contraceptives. *Br J Clin Pharmacol*. 1989;27(6):851-859.
62. Sterling JB, Heymann WR. Potassium iodide in dermatology: a 19th century drug for the 21st century—uses, pharmacology, adverse effects, and contraindications. *J Am Acad Dermatol*. 2000;43(4):691-697.
63. Barbour GL, Coburn JW, Slatopolsky E, Norman AW, Horst RL. Hypercalcemia in an anephric patient with sarcoidosis: evidence for extrarenal generation of 1,25-dihydroxyvitamin D. *N Engl J Med*. 1981;305(8):440-443.
64. Adams JS, Sharma OP, Gacad MA, Singer FR. Metabolism of 25-hydroxyvitamin D3 by cultured pulmonary alveolar macrophages in sarcoidosis. *J Clin Invest*. 1983;72(5):1856-1860.
65. Spindel SJ, Hamill RJ, Georgiou PR, Lacke CE, Green LK, Mallette LE. Case report: vitamin D-mediated hypercalcemia in fungal infections. *Am J Med Sci*. 1995;310(2):71-76.
66. Kantarjian HM, Saad MF, Estey EH, Sellin RV, Samaan NA. Hypercalcemia in disseminated candidiasis. *Am J Med*. 1983;74(4):721-724.
67. Ali MY, Gopal KV, Llerena LA, Taylor HC. Hypercalcemia associated with infection by *Cryptococcus neoformans* and *Coccidioides immitis*. *Am J Med Sci*. 1999;318(6):419-423.
68. Caldwell JW, Arsura EL, Kilgore WB, Reddy CM, Johnson RH. Hypercalcemia in patients with disseminated coccidioidomycosis. *Am J Med Sci*. 2004;327(1):15-18.
69. Silva LC, Ferrari TC. Hypercalcaemia and paracoccidioidomycosis. *Trans R Soc Trop Med Hyg*. 1998;92(2):187.
70. Ahmed B, Jaspan JB. Case report: hypercalcemia in a patient with AIDS and *Pneumocystis carinii* pneumonia. *Am J Med Sci*. 1993;306(5):313-316.
71. Stewart AF, Horst R, Deftos LJ, Cadman EC, Lang R, Broadus AE. Biochemical evaluation of patients with cancer-associated hypercalcemia: evidence for humoral and nonhumoral groups. *N Engl J Med*. 1980;303(24):1377-1383.
72. Jenny-Avital ER, Abadi M. Immune reconstitution cryptococcosis after initiation of successful highly active antiretroviral therapy. *Clin Infect Dis*. 2002 Dec 15;35(12):e128-e133. Epub 2002 Nov 20.
73. Lawn SD, Macallan DC. Hypercalcemia: a manifestation of immune reconstitution complicating tuberculosis in an HIV-infected person [letter]. *Clin Infect Dis*. 2004;38(1):154-155.
74. Reinholt FP, Hultenby K, Oldberg A, Heinegård D. Osteopontin—a possible anchor of osteoclasts to bone. *Proc Natl Acad Sci U S A*. 1990;87(12):4473-4475.
75. Carlson I, Tognazzi K, Manseau EJ, Dvorak HF, Brown LF. Osteopontin is strongly expressed by histiocytes in granulomas of diverse etiology. *Lab Invest*. 1997;77(1):103-108.
76. Li J, Sarosi I, Yan XQ, et al. RANK is the intrinsic hematopoietic cell surface receptor that controls osteoclastogenesis and regulation of bone mass and calcium metabolism. *Proc Natl Acad Sci U S A*. 2000;97(4):1566-1571.
77. Siddiqi A, Goh BT, Brown CL, Hillman RJ, Monson JP. Hypothyroidism and hypoparathyroidism associated with *Pneumocystis carinii* infection in a patient with AIDS. *Int J STD AIDS*. 1998;9(2):108-110.
78. Sonino N. The endocrine effects of ketoconazole. *J Endocrinol Invest*. 1986;9(4):341-347.
79. Henry HL. Effect of ketoconazole and miconazole on 25-hydroxyvitamin D3 metabolism by cultured chick kidney cells. *J Steroid Biochem*. 1985;23(6A):991-994.
80. Glass AR, Eil C. Ketoconazole-induced reduction in serum 1,25-dihydroxyvitamin D and total serum calcium in hypercalcemic patients. *J Clin Endocrinol Metab*. 1988;66(5):934-938.
81. Saggese G, Bertelloni S, Baroncelli GI, Di Nero G. Ketoconazole decreases the serum ionized calcium and 1,25-dihydroxyvitamin D levels in tuberculosis-associated hypercalcemia. *Am J Dis Child*. 1993;147(3):270-273.
82. Adams JS, Sharma OP, Diz MM, Endres DB. Ketoconazole decreases the serum 1,25-dihydroxyvitamin D and calcium concentration in sarcoidosis-associated hypercalcemia. *J Clin Endocrinol Metab*. 1990;70(4):1090-1095.
83. Marcus N, Garty BZ. Transient hypoparathyroidism due to amphotericin B-induced hypomagnesemia in a patient with beta-thalassemia. *Ann Pharmacother*. 2001;35(9):1042-1044.
84. Offner F, Krcmery V, Boogaerts M, et al; EORTC Invasive Fungal Infections Group. Liposomal nystatin in patients with invasive aspergillosis refractory to or intolerant of amphotericin B. *Antimicrob Agents Chemother*. 2004;48(12):4808-4812.
85. Maertens J, Raad I, Petrikos G, et al; Caspofungin Salvage Aspergillosis Study Group. Efficacy and safety of caspofungin for treatment of invasive aspergillosis in patients refractory to or intolerant of conventional antifungal therapy. *Clin Infect Dis*. 2004 Dec 1;39(11):1563-1571. Epub 2004 Nov 9.
86. Jain A, Butani L. Severe hyperphosphatemia resulting from high-dose liposomal amphotericin in a child with leukemia. *J Pediatr Hematol Oncol*. 2003;25(4):324-326.
87. Mendoza D, Connors S, Lane C, Stehna S. Liposomal amphotericin B as a cause of pseudohyperphosphatemia [letter]. *Clin Infect Dis*. 2008;46(4):645-646.
88. Guice KS, Lynch M, Weatherbee L. Invasive aspergillosis: an unusual cause of hemorrhagic pancreatitis. *Am J Gastroenterol*. 1987;82(6):563-565.
89. Parenti DM, Steinberg W, Kang P. Infectious causes of acute pancreatitis. *Pancreas*. 1996;13(4):356-371.
90. Ribes JA, Vanover-Sams CL, Baker DJ. *Zygomycetes* in human disease. *Clin Microbiol Rev*. 2000;13(2):236-301.

91. Richardson SE, Bannatyne RM, Summerbell RC, Milliken J, Gold R, Weitzman SS. Disseminated fusarial infection in the immunocompromised host. *Rev Infect Dis*. 1988;10(6):1171-1181.
92. Rippon JW, Zvetina JR, Reyes C. Case report: miliary blastomycosis with cerebral involvement. *Mycopathologia*. 1977;60(2):121-125.
93. Goodwin RA Jr, Shapiro JL, Thurman GH, Thurman SS, Des Prez RM. Disseminated histoplasmosis: clinical and pathologic correlations. *Medicine (Baltimore)*. 1980;59(1):1-33.
94. Silva-Vergara ML, Maneira FR, De Oliveira RM, Santos CT, Etchebehere RM, Adad SJ. Multifocal sporotrichosis with meningeal involvement in a patient with AIDS. *Med Mycol*. 2005;43(2):187-190.
95. Bigliazzi C, Poletti V, Dell'Amore D, Saragoni L, Colby TV. Disseminated basidiobolomycosis in an immunocompetent woman. *J Clin Microbiol*. 2004;42(3):1367-1369.
96. Tsai CY, Lü YC, Wang LT, Hsu TL, Sung JL. Systemic chromoblastomycosis due to *Hormodendrum dermatitidis* (Kano) Conant: report of the first case in Taiwan. *Am J Clin Pathol*. 1966;46(1):103-114.
97. Pynka M, Wnuk A, Bander D, et al. Disseminated infection with *Saccharomyces kluyveri* in a patient with AIDS. *Infection*. 1998;26(3):184-186.
98. Shek YH, Tucker MC, Viciano AL, Manz HJ, Connor DH. *Malassezia furfur*—disseminated infection in premature infants. *Am J Clin Pathol*. 1989;92(5):595-603.
99. Hoerauf A, Hammer S, Müller-Myhsok B, Rupprecht H. Intra-abdominal *Candida* infection during acute necrotizing pancreatitis has a high prevalence and is associated with increased mortality. *Crit Care Med*. 1998;26(12):2010-2015.
100. De Waele JJ, Vogelaers D, Blot S, Colardyn F. Fungal infections in patients with severe acute pancreatitis and the use of prophylactic therapy. *Clin Infect Dis*. 2003 Jul 15;37(2):208-213. Epub 2003 Jul 9.
101. Curley PJ. Endotoxin, cellular immune dysfunction and acute pancreatitis. *Ann R Coll Surg Engl*. 1996;78(6):531-535.
102. Felk A, Kretschmar M, Albrecht A, et al. *Candida albicans* hyphal formation and the expression of the *Efg1*-regulated proteinases Sap4 to Sap6 are required for the invasion of parenchymal organs. *Infect Immun*. 2002;70(7):3689-3700.
103. Trivedi CD, Pitchumoni CS. Drug-induced pancreatitis: an update. *J Clin Gastroenterol*. 2005;39(8):709-716.
104. Stuecklin-Utsch A, Hasan C, Bode U, Fleischhack G. Pancreatic toxicity after liposomal amphotericin B. *Mycoses*. 2002;45(5-6):170-173.
105. Sato K, Hayashi M, Utsugi M, Ishizuka T, Takagi H, Mori M. Acute pancreatitis in a patient treated with micafungin. *Clin Ther*. 2007;29(7):1468-1473.
106. Vazquez JA, Skiest DJ, Nieto L, et al. A multicenter randomized trial evaluating posaconazole versus fluconazole for the treatment of oropharyngeal candidiasis in subjects with HIV/AIDS. *Clin Infect Dis*. 2006;42(8):1179-1186.
107. Boyd AE, Modi S, Howard SJ, Moore CB, Keevil BG, Denning DW. Adverse reactions to voriconazole. *Clin Infect Dis*. 2004;39(8):1241-1244.
108. Balian JD, Sukhova N, Harris JW, et al. The hydroxylation of omeprazole correlates with S-mephenytoin metabolism: a population study. *Clin Pharmacol Ther*. 1995;57(6):662-669.
109. Lobo BL, Miwa LJ, Jungnickel PW. Possible ketoconazole-induced hypoglycemia [letter]. *Drug Intell Clin Pharm*. 1988;22(7-8):632.
110. Niemi M, Backman JT, Neuvonen M, Laitila J, Neuvonen PJ, Kivistö KT. Effects of fluconazole and fluvonoxamine on the pharmacokinetics and pharmacodynamics of glimepiride. *Clin Pharmacol Ther*. 2001;69(4):194-200.
111. Lazar JD, Wilner KD. Drug interactions with fluconazole. *Rev Infect Dis*. 1990;12(suppl 3):S327-S333.
112. Niemi M, Neuvonen M, Juntti-Patinen L, Backman JT, Neuvonen PJ. Effect of fluconazole on the pharmacokinetics and pharmacodynamics of nateglinide. *Clin Pharmacol Ther*. 2003;74(1):25-31.
113. Rowe BR, Thorpe J, Barnett A. Safety of fluconazole in women taking oral hypoglycaemic agents [letter]. *Lancet*. 1992;339(8787):255-256.
114. Miners JO, Birkett DJ. Cytochrome P4502C9: an enzyme of major importance in human drug metabolism. *Br J Clin Pharmacol*. 1998;45(6):525-538.
115. Frenkel JK. Pathogenesis of infections of the adrenal gland leading to Addison's disease in man: the role of corticoids in adrenal and generalized infection. *Ann N Y Acad Sci*. 1960;84:391-440.
116. Lio S, Cibini M, Marcello R, Viviani MA, Ajello L. Adrenal bilateral incidentaloma by reactivated histoplasmosis. *J Endocrinol Invest*. 2000;23(7):476-479.
117. Papadopoulos KI, Castor B, Klingspor L, Dejmeck A, Lorén I, Bramnert M. Bilateral isolated adrenal coccidioidomycosis. *J Intern Med*. 1996;239(3):275-278.
118. Torres CM, Duarte E, Guimaraes JP, Moreira LF. Destructive lesion of the adrenal gland in South American blastomycosis (Lutz' disease). *Am J Pathol*. 1952;28(1):145-155.
119. Agarwal J, Agarwal G, Ayyagari A, Kar DK, Mishra SK, Bhatia E. Isolated *Pneumocystis carinii* infection of adrenal glands causing Addison's disease in a non-immunocompromised adult. *Endocr Pathol*. 2001 Spring;12(1):87-91.
120. Maloney PJ. Addison's disease due to chronic disseminated coccidioidomycosis. *AMA Arch Intern Med*. 1952;90(6):869-878.
121. Del Negro G, Wajchenberg BL, Pereira VG, et al. Addison's disease associated with South American blastomycosis. *Ann Intern Med*. 1961;54:189-197.
122. Shah B, Taylor HC, Pillay I, Chung-Park M, Dobrinich R. Adrenal insufficiency due to cryptococcosis. *JAMA*. 1986;256(23):3247-3249.
123. Kent DC, Collier TM. Addison's disease associated with North American blastomycosis: a case report. *J Clin Endocrinol Metab*. 1965;25:164-169.
124. Zhu QZ, Hu J, Mulay S, Esch F, Shimasaki S, Solomon S. Isolation and structure of corticosteroid peptides from rabbit fetal and adult lung. *Proc Natl Acad Sci U S A*. 1988;85(2):592-596.
125. Alteras I, Cojocaru I, Balanescu A. Generalized candidiasis associated with Addison's disease. *Mykosen*. 1969;12(9):575-577.
126. Karim M, Sheikh H, Alam M, Sheikh Y. Disseminated *Bipolaris* infection in an asthmatic patient: case report. *Clin Infect Dis*. 1993;17(2):248-253.
127. Washburn RG, Bennett JE. Reversal of adrenal glucocorticoid dysfunction in a patient with disseminated histoplasmosis. *Ann Intern Med*. 1989;110(1):86-87.
128. Osa SR, Peterson RE, Roberts RB. Recovery of adrenal reserve following treatment of disseminated South American blastomycosis. *Am J Med*. 1981;71(2):298-301.
129. Del Negro G, Melo EH, Rodbard D, Melo MR, Layton J, Wachslight-Rodbard H. Limited adrenal reserve in paracoccidioidomycosis: cortisol and aldosterone responses to 1-24 ACTH. *Clin Endocrinol (Oxf)*. 1980;13(6):553-559.
130. Leal AM, Magalhães PK, Martinez R, Moreira AC. Adrenocortical hormones and interleukin patterns in paracoccidioidomycosis. *J Infect Dis*. 2003 Jan 1;187(1):124-127. Epub 2002 Dec 13.
131. Kawashima A, Sandler CM, Fishman EK, et al. Spectrum of CT findings in nonmalignant disease of the adrenal gland. *Radiographics*. 1998;18(2):393-412.
132. Lamberts SW, Bons EG, Bruining HA, de Jong FH. Differential effects of the imidazole derivatives etomidate, ketoconazole and miconazole and of metyrapone on the secretion of cortisol and its precursors by human adrenocortical cells. *J Pharmacol Exp Ther*. 1987;240(1):259-264.
133. Loose DS, Kan PB, Hirst MA, Marcus RA, Feldman D. Ketoconazole blocks adrenal steroidogenesis by inhibiting cytochrome P450-dependent enzymes. *J Clin Invest*. 1983;71(5):1495-1499.
134. Sonino N. The use of ketoconazole as an inhibitor of steroid production. *N Engl J Med*. 1987;317(13):812-818.
135. Leal-Cerro A, García Luna PP, Villar J, et al. Ketoconazole as an inhibitor of steroid production [letter]. *N Engl J Med*. 1988;318(11):710-711.
136. Pont A, Williams PL, Loose DS, et al. Ketoconazole blocks adrenal steroid synthesis. *Ann Intern Med*. 1982;97(3):370-372.
137. Pont A, Graybill JR, Craven PC, et al. High-dose ketoconazole therapy and adrenal and testicular function in humans. *Arch Intern Med*. 1984;144(11):2150-2153.
138. Tucker WS Jr, Snell BB, Island DP, Gregg CR. Reversible adrenal insufficiency induced by ketoconazole. *JAMA*. 1985;253(16):2413-2414.
139. Best TR, Jenkins JK, Murphy FY, Nicks SA, Bussell KL, Vesely DL. Persistent adrenal insufficiency secondary to low-dose ketoconazole therapy. *Am J Med*. 1987;82(3):676-680.
140. Loose DS, Stover EP, Feldman D. Ketoconazole binds to glucocorticoid receptors and exhibits glucocorticoid antagonist activity in cultured cells. *J Clin Invest*. 1983;72(1):404-408.
141. Sonino N, Boscaro M, Paoletta A, Mantero F, Ziliotto D. Ketoconazole treatment in Cushing's syndrome: experience in 34 patients. *Clin Endocrinol (Oxf)*. 1991;35(4):347-352.
142. Contreras P, Rojas A, Biagini L, González P, Massardo T. Regression of metastatic adrenal carcinoma during palliative ketoconazole treatment [letter]. *Lancet*. 1985;2(8447):151-152.

143. Shepherd FA, Hoffert B, Evans WK, Emery G, Trachtenberg J. Ketoconazole: use in the treatment of ectopic adrenocorticotrophic hormone production and Cushing's syndrome in small-cell lung cancer. *Arch Intern Med*. 1985;145(5):863-864.
144. Duret C, Daujat-Chavanier M, Pascussi JM, et al. Ketoconazole and miconazole are antagonists of the human glucocorticoid receptor: consequences on the expression and function of the constitutive androstane receptor and the pregnane X receptor. *Mol Pharmacol*. 2006 Jul;70(1):329-339. Epub 2006 Apr 11.
145. Benito P, Corpas MS, Quesada JM, Jiménez JA. Control of hyperaldosteronism by ketoconazole [letter]. *Drug Intell Clin Pharm*. 1987; 21(9):752.
146. Aabo K, De Coster R. Hypertension during high-dose ketoconazole treatment: a probable mineralocorticosteroid effect [letter]. *Lancet*. 1987;2(8559):637-638.
147. Albert SG, DeLeon MJ, Silverberg AB. Possible association between high-dose fluconazole and adrenal insufficiency in critically ill patients. *Crit Care Med*. 2001;29(3):668-670.
148. Magill SS, Puthanakit T, Swoboda SM, et al. Impact of fluconazole prophylaxis on cortisol levels in critically ill surgical patients. *Antimicrob Agents Chemother*. 2004;48(7):2471-2476.
149. Santhana Krishnan SG, Cobbs RK. Reversible acute adrenal insufficiency caused by fluconazole in a critically ill patient. *Postgrad Med J*. 2006;82(971):e23.
150. Riedl M, Maier C, Zetting G, Nowotny P, Schima W, Luger A. Long term control of hypercortisolism with fluconazole: case report and in vitro studies. *Eur J Endocrinol*. 2006;154(4):519-524.
151. Queiroz-Telles F, Purim KS, Boguszewski CL, Afonso FC, Graf H. Adrenal response to corticotrophin and testosterone during long-term therapy with itraconazole in patients with chromoblastomycosis. *J Antimicrob Chemother*. 1997;40(6):899-902.
152. Sharkey PK, Rinaldi MG, Dunn JF, Hardin TC, Fetchick RJ, Graybill JR. High-dose itraconazole in the treatment of severe mycoses. *Antimicrob Agents Chemother*. 1991;35(4):707-713.
153. Skov M, Main KM, Sillesen IB, Müller J, Koch C, Lanng S. Iatrogenic adrenal insufficiency as a side-effect of combined treatment of itraconazole and budesonide. *Eur Respir J*. 2002;20(1):127-133.
154. Varis T, Kivistö KT, Backman JT, Neuvonen PJ. Itraconazole decreases the clearance and enhances the effects of intravenously administered methylprednisolone in healthy volunteers. *Pharmacol Toxicol*. 1999;85(1):29-32.
155. DeRijk RH, Schaaf M, de Kloet ER. Glucocorticoid receptor variants: clinical implications. *J Steroid Biochem Mol Biol*. 2002;81(2):103-122.
156. Deray G. Amphotericin B nephrotoxicity. *J Antimicrob Chemother*. 2002;49(suppl 1):37-41.
157. Mohan S, Ahmed S, Alimohammadi B, Jaitly M, Cheng JT, Pogue VA. Proteinuria lowers the risk of amphotericin B-associated hypokalaemia. *J Antimicrob Chemother*. 2007 Sep;60(3):690-693. Epub 2007 Jun 27.
158. Cornely OA, Maertens J, Bresnik M, et al; AmBiLoad Trial Study Group. Liposomal amphotericin B as initial therapy for invasive mold infection: a randomized trial comparing a high-loading dose regimen with standard dosing (AmBiLoad Trial). *Clin Infect Dis*. 2007 May 15;44(10):1289-1297. Epub 2007 Apr 9.
159. Wazny LD, Brophy DF. Amiloride for the prevention of amphotericin B-induced hypokalemia and hypomagnesemia. *Ann Pharmacother*. 2000; 34(1):94-97.
160. Ural AU, Avcu F, Cetin T, et al. Spironolactone: is it a novel drug for the prevention of amphotericin B-related hypokalemia in cancer patients? *Eur J Clin Pharmacol*. 2002;57(11):771-773.
161. Nucci M, Loureiro M, Silveira F, et al. Comparison of the toxicity of amphotericin B in 5% dextrose with that of amphotericin B in fat emulsion in a randomized trial with cancer patients. *Antimicrob Agents Chemother*. 1999; 43(6):1445-1448.
162. Glasmacher A, Cornely O, Ullmann AJ, et al; Itraconazole Research Group of Germany. An open-label randomized trial comparing itraconazole oral solution with fluconazole oral solution for primary prophylaxis of fungal infections in patients with haematological malignancy and profound neutropenia. *J Antimicrob Chemother*. 2006 Feb;57(2):317-325. Epub 2005 Dec 8.
163. Denning DW, Ribaud P, Milpied N, et al. Efficacy and safety of voriconazole in the treatment of acute invasive aspergillosis. *Clin Infect Dis*. 2002 Mar 1;34(5):563-571. Epub 2002 Jan 22.
164. Nelson MR, Smith D, Erskine D, Gazzard BG. Ventricular fibrillation secondary to itraconazole induced hypokalaemia [letter]. *J Infect*. 1993;26(3):348.
165. Arathoon EG, Gotuzzo E, Noriega LM, Berman RS, DiNubile MJ, Sable CA. Randomized, double-blind, multicenter study of caspofungin versus amphotericin B for treatment of oropharyngeal and esophageal candidiasis. *Antimicrob Agents Chemother*. 2002;46(2):451-457.
166. Craven PC, Gremillion DH. Risk factors of ventricular fibrillation during rapid amphotericin B infusion. *Antimicrob Agents Chemother*. 1985; 27(5):868-871.
167. Barton CH, Pahl M, Vaziri ND, Cesario T. Renal magnesium wasting associated with amphotericin B therapy. *Am J Med*. 1984;77(3):471-474.
168. Kraemer FB, Pont A. Inhibition of cholesterol synthesis by ketoconazole. *Am J Med*. 1986;80(4):616-622.
169. Dushkin MI, Zenkov NK, Menshikova EB, Pivovarov EN, Lyubimov GY, Volkov NN. Ketoconazole inhibits oxidative modification of low density lipoprotein. *Atherosclerosis*. 1995;114(1):9-18.
170. Kesäniemi YA, Miettinen TA. Inhibition of cholesterol absorption by neomycin, benzodiazepine derivatives and ketoconazole. *Eur J Clin Pharmacol*. 1991;40(suppl 1):S65-S67.
171. Gylling H, Vanhanen H, Miettinen TA. Effects of ketoconazole on cholesterol precursors and low density lipoprotein kinetics in hypercholesterolemia. *J Lipid Res*. 1993;34(1):59-67.
172. Dismukes WE, Stamm AM, Graybill JR, et al. Treatment of systemic mycoses with ketoconazole: emphasis on toxicity and clinical response in 52 patients: National Institute of Allergy and Infectious Diseases collaborative antifungal study. *Ann Intern Med*. 1983;98(1):13-20.
173. Tucker RM, Haq Y, Denning DW, Stevens DA. Adverse events associated with itraconazole in 189 patients on chronic therapy. *J Antimicrob Chemother*. 1990;26(4):561-566.
174. Wise GJ, Talluri GS, Marella VK. Fungal infections of the genitourinary system: manifestations, diagnosis, and treatment. *Urol Clin North Am*. 1999;26(4):701-718.
175. Powell CR, Allshouse M, Bethel KJ, Mevorach RA. Invasive aspergillosis of the scrotum. *J Urol*. 1998;159(4):1306-1308.
176. Tomimori-Yamashita J, Tagliolatto S, Porro AM, Ogawa MM, Michalany NS, Camargo ZP. Paracoccidioidomycosis: an uncommon localization in the scrotum. *Mycoses*. 1997;40(11-12):415-418.
177. James CL, Lomax-Smith JD. Cryptococcal epididymo-orchitis complicating steroid therapy for relapsing polychondritis. *Pathology*. 1991;23(3): 256-258.
178. Hironaga M, Okazaki N, Saito K, Watanabe S. *Trichophyton mentagrophytes* granulomas: unique systemic dissemination to lymph nodes, testes, vertebrae, and brain. *Arch Dermatol*. 1983;119(6):482-490.
179. Patel PJ, Pareek SS. Scrotal ultrasound in male infertility. *Eur Urol*. 1989;16(6):423-425.
180. Nagy B, Sutka P. Demonstration of antibodies against *Candida guilliermondii* var. *guilliermondii* in asymptomatic infertile men. *Mycoses*. 1992;35(9-10):247-250.
181. Nagy B, Sutka P, Ziwe-el-Abidine M, et al. *Candida guilliermondii* var. *guilliermondii* infection in infertile women. *Mycoses*. 1989;32(9):463-468.
182. Tuttle JP Jr, Bannister ER, Derrick FC. Interference of human spermatozoal motility and spermatozoal agglutination by *Candida albicans*. *J Urol*. 1977;118(5):797-799.
183. Staib F, Seibold M, L'age M, et al. *Cryptococcus neoformans* in the seminal fluid of an AIDS patient: a contribution to the clinical course of cryptococcosis. *Mycoses*. 1989;32(4):171-180.
184. Zinedine A, Soriano JM, Moltó JC, Mañes J. Review on the toxicity, occurrence, metabolism, detoxification, regulations and intake of zearalenone: an oestrogenic mycotoxin. *Food Chem Toxicol*. 2007 Jan;45(1):1-18. Epub 2006 Aug 30.
185. Schürmeyer T, Nieschlag E. Effect of ketoconazole and other imidazole fungicides on testosterone biosynthesis. *Acta Endocrinol (Copenh)*. 1984;105(2):275-280.
186. Pont A, Williams PL, Azhar S, et al. Ketoconazole blocks testosterone synthesis. *Arch Intern Med*. 1982;142(12):2137-2140.
187. Pont A, Goldman ES, Sugar AM, Siiteri PK, Stevens DA. Ketoconazole-induced increase in estradiol-testosterone ratio: probable explanation for gynecomastia. *Arch Intern Med*. 1985;145(8):1429-1431.
188. Glass AR. Ketoconazole-induced stimulation of gonadotropin output in men: basis for a potential test of gonadotropin reserve. *J Clin Endocrinol Metab*. 1986;63(5):1121-1125.

189. Trachtenberg J, Pont A. Ketoconazole therapy for advanced prostate cancer. *Lancet*. 1984;2(8400):433-435.
190. Holland FJ, Fishman L, Bailey JD, Fazekas AT. Ketoconazole in the management of precocious puberty not responsive to LHRH-analogue therapy. *N Engl J Med*. 1985;312(16):1023-1028.
191. Hugo Perez BS. Ketoconazole as an adjunct to finasteride in the treatment of androgenetic alopecia in men. *Med Hypotheses*. 2004;62(1):112-115.
192. Faredin I, Tóth I, Oszlányi J, Scultéry S. In vitro study of rat prostate 5 alpha-reductase activity and its inhibition. *Int Urol Nephrol*. 1992;24(2):145-154.
193. Eil C. Ketoconazole binds to the human androgen receptor. *Horm Metab Res*. 1992;24(8):367-370.
194. el-Medany AH, Hagar HH. Effect of fluconazole on the fertility of male rabbits. *Arzneimittelforschung*. 2002;52(8):636-640.
195. Hanger DP, Jevons S, Shaw JT. Fluconazole and testosterone: in vivo and in vitro studies. *Antimicrob Agents Chemother*. 1988;32(5):646-648.
196. Foote RH. Spermicidal effects of amphotericin B and nystatin on bull and rabbit sperm and contraceptive effects in rabbits. *Contraception*. 2002;66(3):193-197.
197. Kim SW, Nah MY, Yeum CH, et al. Pelvic aspergillosis with tubo-ovarian abscess in a renal transplant recipient. *J Infect*. 2001;42(3):215-217.
198. Rodriguez M, Okagaki T, Richart RM. Mycotic endometritis due to *Candida*: a case report. *Obstet Gynecol*. 1972;39(2):292-294.
199. Craig MW, Davey WN, Green RA. Conjugal blastomycosis. *Am Rev Respir Dis*. 1970;102(1):86-90.
200. Chowfin A, Tight R. Female genital coccidioidomycosis (FGC), Addison's disease and sigmoid loop abscess due to *Coccidioides immitis*: case report and review of literature on FGC. *Mycopathologia*. 1999;145(3):121-126.
201. Sheyn I, Mira JL, Thompson MB. *Paracoccidioides brasiliensis* in a postpartum Pap smear: a case report. *Acta Cytol*. 2001;45(1):79-81.
202. Sills M, Schwartz A, Weg JG. Conjugal histoplasmosis: a consequence of progressive dissemination in the index case after steroid therapy. *Ann Intern Med*. 1973;79(2):221-224.
203. Padhye AA, Hall L, Roberts GD. Deep-seated trichosporonosis in an immunocompetent patient: a case report of uterine trichosporonosis-correction [letter]. *Clin Infect Dis*. 2002;35(1):108-109.
204. Arora NS, Oblinger MJ, Feldman PS. Chronic pleural blastomycosis with hyperprolactinemia, galactorrhea, and amenorrhea. *Am Rev Respir Dis*. 1979;120(2):451-455.
205. Weber MM, Will A, Adelman B, Engelhardt D. Effect of ketoconazole on human ovarian C17,20-desmolase and aromatase. *J Steroid Biochem Mol Biol*. 1991;38(2):213-218.
206. Harris AL, Cantwell BM, Dowsett M. High dose ketoconazole: endocrine and therapeutic effects in postmenopausal breast cancer. *Br J Cancer*. 1988;58(4):493-496.
207. Cummings AM, Hedge JL, Laskey J. Ketoconazole impairs early pregnancy and the decidual cell response via alterations in ovarian function. *Fundam Appl Toxicol*. 1997;40(2):238-246.
208. National Institutes of Allergy and Infectious Diseases Mycoses Study Group. Treatment of blastomycosis and histoplasmosis with ketoconazole: results of a prospective randomized clinical trial. *Ann Intern Med*. 1985;103(6, pt 1):861-872.
209. Vidal-Puig AJ, Muñoz-Torres M, Jódar-Gimeno E, et al. Ketoconazole therapy: hormonal and clinical effects in non-tumoral hyperandrogenism. *Eur J Endocrinol*. 1994;130(4):333-338.
210. Raad II, Graybill JR, Bustamante AB, et al. Safety of long-term oral posaconazole use in the treatment of refractory invasive fungal infections. *Clin Infect Dis*. 2006 Jun 15;42(12):1726-1734. Epub 2006 May 8.
211. Sinofsky FE, Pasquale SA. The effect of fluconazole on circulating ethinyl estradiol levels in women taking oral contraceptives. *Am J Obstet Gynecol*. 1998;178(2):300-304.
212. Andrews E, Damle BD, Fang A, et al. Pharmacokinetics and tolerability of voriconazole and a combination oral contraceptive co-administered in healthy female subjects. *Br J Clin Pharmacol*. 2008 Apr;65(4):531-539. Epub Feb 21.
213. Pillans PI, Sparrow MJ. Pregnancy associated with a combined oral contraceptive and itraconazole [letter]. *N Z Med J*. 1993;106(965):436.